

Reactivation of *Serratia marcescens* mutant endonuclease by Hydroxylamine

Khamidullina R., Fazleeva I., Trushin M., Gimadutdinov O.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018, Pharmainfo Publications. All rights reserved. It is known that histidine plays an important role in the catalytic activity of many nucleases. Performing the function of a common base of these enzymes, it activates the formation of hydroxyl from the water molecule, which, in turn, by attacking the phosphorus atom of the diester bond causes its rupture. It was previously shown that in the endonuclease *Serratia marcescens*, the replacement of histidine with glycine results in its inactivation. We were able to restore the hydroxylamine activity of the mutant enzyme *Serratia marcescens* endonuclease, in which histidine in the 89th position is replaced by glycine.

Keywords

Basal level of expression, Endonuclease, Histidine, Hydroxylamine, Plasmid

References

- [1] Leshchinskaya, I.B. A comparative study of bacterial phosphomonosomes phosphodiesterases hydrolyzing nucleic acids and nucleotides. Leshchinskaya, M.I. Belyaeva, K.Z. Gilfanov// *Microbiology*.-1968.-P. 37.-p. 979-983.
- [2] Gimadutdinov, O.A. Synthesis of extracellular endonuclease *Serratia marcescens* recombinant strains of *Escherichia coli*. Gimadutdinov, L.M. Antsilevich// *Biotechnology*.-1993.-№5.
- [3] Friedhoff P. A procedure for renaturation and purification of the extracellular *Serratia marcescens* nuclease from genetically engineered *Escherichia coli*/ P. Friedhoff, O. Gimadutdinov, T. Ruter, W. Wende, C. Urbanke, H. Thole, A. Pingoud// *Protein Expr.Purif.* - 1994a. - Vol.5. - p. 37-43.
- [4] Friedhoff P. Identification of catalytically relevant amino acids of the extracellular *Serratia marcescens* endonuclease by alignment-guided mutagenesis/ P. Friedhoff, O. Gimadutdinov, A. Pingoud// *Nucleic Acids Research*. - 1994. - Vol. 22, No. 16. - 3280 - 3287.
- [5] Friedhoff, P. Analysis of the mechanism of the *Serratia* nuclease using site-directed mutagenesis/ P. Friedhoff, B. Kolmes, O. Gimadutdinov, W. Wende, K.L. Krause, A. Pingoud// *Nucleic Acids Reserch*. - 1996a. - Vol. 24. - p. 2632 - 2639.
- [6] Venkataraman P. Chemical rescue of histidine selectivity filter mutants of the M2 ion channel of influenza A virus/ P. Venkataraman, R.A.Lamb, L.H.Pinto// *J. Biological Chemistry*. - 2005. - Vol. 280, No.22, -P.21463-21472.
- [7] Peracchi A. How (and why) to revive a dead enzyme: the power of chemical rescue/ A. Peracchi// *Current Chem. Biol.* - 2008. - Vol. 2, No. 1. - P 32 - 49.
- [8] Chen W.J. Probing the catalytic mechanism of bovine pancreatic deoxyribonuclease I by chemical rescue/ W.J. Chen, P.J. Lai, Y.S. Lai, P.T. Huang, C.C. Lin, T.H. Liao// *Biochemical and Biophysical Research Communications* - 2007. - Vol. 352. - P. 689 - 696.

- [9] Midon M. Chemical rescue of active site mutants of nucleases of the H-N-H family by imidazole/ M. Midon, O. Gimadutdinow, G. Meiss, P. Friedhoff, A. Pingoud// *Chembiochem.* – 2012. – Vol. 13, No. 5. – P. 713 – 721.
- [10] Meiss G. Microtiter-plate assay for nonspecific endonucleases [Text]/ G. Meiss, O. Gimadutdinow, P. Friedhoff, A. Pingoud// *Methods Mol Biol.* – 2001. – Vol. 160. – P. 37 – 48.